











NWT Immunization Schedule

Publicly funded vaccines for infants and children

- This is the recommended immunization schedule for those aged birth to 17 years.
- It is never too late to start or catch up on immunizations.
- Delayed or missed vaccines may make it necessary to follow a different schedule.
- If you have any questions, contact your health centre or public health unit.

JUNE 2017

AGE	VACCINE
 Birth	HB Hepatitis B
	BCG¹ Bacillus Calmette–Guérin (Tuberculosis)
1 month	HB Hepatitis B
 2 months	DTaP-IPV-Hib Diphtheria, tetanus, acellular pertussis, inactivated polio, <i>Haemophilus influenzae</i> type B
	Men-C-C Meningococcal conjugate C
	Pneu-C-13 Pneumococcal conjugate 13
	Rot-1 Rotavirus
4 months	DTaP-IPV-Hib Diphtheria, tetanus, acellular pertussis, inactivated polio, <i>Haemophilus influenzae</i> type B
	Pneu-C-13 Pneumococcal conjugate 13
	Rot-1 Rotavirus
 6 months	DTaP-IPV-Hib Diphtheria, tetanus, acellular pertussis, inactivated polio, <i>Haemophilus influenzae</i> type B
	HB Hepatitis B
	Pneu-C-13 Pneumococcal conjugate 13
12 months	MMRV Measles, mumps, rubella, varicella
	Men-C-C Meningococcal conjugate C
 18 months	DTaP-IPV-Hib Diphtheria, tetanus, acellular pertussis, inactivated polio, <i>Haemophilus influenzae</i> type B
	Pneu-C-13 Pneumococcal conjugate 13
3 years	MMRV Measles, mumps, rubella, varicella
4-6 years	DTaP-IPV² Diphtheria, tetanus, acellular pertussis and inactivated polio
 Grades 4, 5 or 6	HPV Human papillomavirus
	9 to 14 years – 2 dose series 15 years and older – 3 dose series
 Grade 7	Tdap Tetanus, diphtheria, acellular pertussis
 Grade 12	Men-C-ACYW-135³ Meningococcal conjugate quadrivalent
 6 months and older	Inf Influenza (flu) offered yearly. Children 8 years of age and younger receiving this vaccine for the first time will need a 2nd dose 1 month after the 1st dose.

1. BCG is only recommended for certain infants. Talk to your health care provider or public health nurse for more information.

2. DTaP-IPV-Hib may be substituted for DTaP-IPV in times of shortage.

3. Vaccine offered only to those grade 12 students who will be attending post-secondary education outside of the NWT.

Publicly funded vaccines for adults

- Immunizations are not just for kids. Adults need vaccines too.
- These are the recommended vaccines for adults and may be dependent on age.
- Depending on the vaccine, more than one dose may be needed.

VACCINE	RECOMMENDATION
HB Hepatitis B	For those who have not received it in childhood
HPV Human papillomavirus	Females and males up to 26 years of age
Inf Influenza	Yearly for everyone 6 months of age and older
MMR Measles, mumps, rubella	For adults born on or after January 1, 1970
Pneu-P-23 Pneumococcal polysaccharide 23	<ul style="list-style-type: none"> • 65 years of age and older • 65 years of age and younger who are at risk for invasive pneumococcal disease (talk to your health care provider)
Tdap Tetanus, diphtheria, pertussis	Every 10 years for everyone
Var Varicella	For those who have never had chickenpox disease

Non-publicly funded vaccines for infants, children and adults

- In addition to the publicly funded vaccines in the NWT, other vaccines are licensed in Canada and are available at a cost.
- Depending on health status some of these vaccines may be provided for free.
- Talk to your health care provider or public health nurse to see if any of these vaccines may be right for you or your child.

There may be a charge for these vaccines

VACCINE	RECOMMENDATION
4CMenB Meningococcal B	Those 2 months to 17 years of age
HPV Human papillomavirus	Females 27 to 45 years of age *vaccine not yet licensed for use in males in this age range
Zos Herpes zoster (shingles)	Adults 50 years of age and older
Pneu-C-13 Pneumococcal conjugate 13	For those 5 years of age and older with certain health conditions

Book travel health appointments at least 6-8 weeks before date of departure

Vaccines for Occupational Health & Safety or Travel

- Other vaccines may be recommended or required depending on the type of work or travel destination. These may include vaccines against: hepatitis A, typhoid, meningococcal disease, Japanese encephalitis, anthrax, rabies and yellow fever.
- Talk to your employer to find out what vaccines may be recommended or required for your type of work.
- Talk to the nurse in your health center or public health clinic to find out what vaccines may be recommended or required for travel outside of Canada.

If you would like this information in another official language, contact us at 1-855-846-9601.

Si vous voulez ces informations dans une autre langue officielle, téléphonez-nous au 1-855-846-9601.