



Last name: _____		First Name: _____	
DOB: _____		Name Alert ☐	
Community: _____			
HCN: _____			
Allergies Known: _____			

Medication Order	(DOPT ONLY)
Weight:	☐ kg ☐ lbs
Dose of INH:	
Dose of Rifapentine:	
Dosage Vitamin B6:	
Ordering Practitioner:	

To be eligible for the 3HP regimen (12 dose, once-weekly combination of isoniazid and rifapentine), patients must have a diagnosis of TB infection (TBI), have TB disease (TBD) ruled out and attend appointments in person for assessment and medication administration (DOPT). Preventative 3HP regimen is the preferred first-line treatment for treatment of TBI and is approved for individuals aged 2 years and older. Eligibility for treatment in children should be assessed by a pediatrician.

Treatment protocol: Must complete 12 doses in 16 weeks to be considered effective treatment.

[illegible]

Common side effects – dose does not need to be withheld. If symptoms persist, consult with MRP												
Nausea or stomach upset (offer food with each dose)	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Loss of appetite	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Fatigue or weakness	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Mild headache	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Muscle aches	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Orange-red discoloration of urine, tears, or other body fluids (expected with rifapentine).	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Withhold dose if ANY of the following symptoms are present and report promptly to ordering practitioner (symptoms may reflect developing hypersensitivity rxn)												
Angioedema, anaphylaxis, Rash, hives, itching	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Hypotension (BP <90/60), dizziness or fainting	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Shortness of breath or wheezing	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Fever or flu-like symptoms	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Numbness or tingling in the hands or feet	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐

HCN:

Persistent nausea, vomiting, diarrhea, or abdominal pain	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Severe headache	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Withhold dose if ANY of the following symptoms are present and report promptly to ordering practitioner (symptoms may reflect potential liver injury or toxicity)												
Yellowing of the skin or eyes (jaundice)	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Dark/brown urine	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Pale stools	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Severe fatigue	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Persistent loss of appetite	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Abdominal pain, especially on the right side	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Medication administration record (Initial with date and time if dose administered. Indicate if dose was HELD. ***Use BCCDC 3HP regimen for dosing reference below).												
Rifapentine ____mg po once weekly	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Isoniazid ____mg po once weekly	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Vitamin B6 ____mg po once weekly	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐	☐
Comments:												

- For patients taking **3HP**, the appearance of **anaphylaxis/angioedema, respiratory symptoms, hypotension, loss of consciousness, jaundice, significant bleeding, or severe systemic symptoms** warrants urgent medical assessment and typically temporary discontinuation of therapy pending evaluation from MRP.
- *Baseline laboratory investigations should be completed as per the NWT TB assessment form Section 3 and reviewed prior to first dose. Ideally preventative therapy should be implemented within one month of completion of baseline investigation/labs. Recommendations for 3HP monitoring and laboratory investigations must be provided by the ordering provider based on health status, side effects, or adverse reactions experienced by the client.
- **Symptom inquiry – **MUST ASK ALL at EVERY VISIT prior to every dose administered.** Patients may experience symptoms after their last dose that resolve prior to their next dose. Any symptoms experienced, whether present now or those that occurred after the last dose, must indicate a YES by “✓” on the form. Provide details of symptom onset, resolution, severity, etc in EMR encounter and NOTIFY ordering practitioner immediately as indicated.
- ***See the [BCCDC Communicable Disease Control Manual Chapter 4: Tuberculosis Section 6: TB Preventive Treatment Table 6-2: Summary of drugs and dosing for TB preventive treatment regimens \(6-2\)](#) Column Once-Weekly Combination of Rifapentine and Isoniazid