

# NWT Patient Teaching Tool

## Sexually Transmitted and Blood-borne Infections (STBBI)



### HOW DO YOU GET CHLAMYDIA/GONORRHEA?



Vaginal Sex



Anal Sex



Oral Sex



Vertical Transmission  
(Mother to baby)

### HOW DO YOU GET SYPHILIS?



Vaginal Sex



Anal Sex



Oral Sex



Vertical Transmission  
(Mother to baby)

### HOW DO YOU GET HIV?



Injection Drug Use



Vaginal Sex



Anal Sex



Vertical Transmission  
(Mother to baby)

### HOW DO YOU GET HEPATITIS C?



Injection Drug Use  
and Sharing Needles



Sharing Smoking Paraphernalia



Tattooing<sup>1</sup>

### RISK FACTORS THAT INCREASE THE POSSIBILITY OF A SEXUALLY TRANSMITTED OR BLOOD BORNE INFECTION:

- Oral, anal or genital sex without a condom/dental dam.
- Multiple sexual partners.
- Sharing injection drug use and non-injection drug use equipment (syringes and pipes).
- <sup>1</sup>Tattoos or piercings with a needle that is not new or sterilized.
- Exchange or sharing of intimate or personal hygiene items (razor, toothbrush, sex toy) that may have been in contact with infected body fluids or blood.