



Instructions for Sexually Transmitted Infections Case Investigation Form

Healthcare professionals (HCPs) have a legislative duty to report notifiable and reportable diseases to the Chief Public Health Officer (CPHO) as required in the [NWT Public Health Act](#) (2009, SNWT 2007, c.17, SI-007-2009), with required information and timelines for reporting as defined in the [Disease Surveillance Regulations](#) (2009, R-096-2009). This information is used for territorial and national surveillance and informs public health planning and interventions. This reporting by the HCPs is accomplished by submitting the [Sexually Transmitted Infections \(STI\) case investigation and contact tracing form](#) to the Office of the Chief Public Health Officer (OCPHO). Information on cases of the following diseases is reportable within specific time frames:

STI	Timeline for submitting <i>Case Investigation Form</i> to OCPHO after making a diagnosis or opinion	Sections of <i>Case Investigation Form</i> to complete
Chlamydial infection	24 hours	All (Sections 1-6) and contact tracing form
Gonococcal infection	24 hours	All (Sections 1-6) and contact tracing form
Syphilis	24 hours	All (Sections 1-6) and contact tracing form
HIV infection	24 hours	Sections 1, 2, & 6 and contact tracing form; contact OCPHO for additional reporting requirements

In addition to case information, HCPs shall make reasonable efforts to initiate contact tracing within 24 hours of reportable disease diagnosis and provide the OCPHO with information respecting the contact tracing and specific control measures that have been initiated or carried out, as outlined in the [Reportable Disease Control Regulations](#) (R-128-2009).

What to Report

Case information (pages 1-2 of [STI case investigation and contact tracing form](#)):

- Sections 1 to 3: Case's confirmed address and telephone number, reason for healthcare visit, and results of physical exam.
 - Physical exam findings help practitioners to stage syphilis infections. For example, if a newly diagnosed syphilis case presents with a rash on their palms or soles of the feet, the infection is likely in its secondary stage¹. Information from this section can also be analyzed for surveillance purposes.
- Sections 4 and 5: Specimen collection information and treatment administered
- Section 6: Risk factors
 - Information in this section is used for surveillance. It can help track the progression of an outbreak through different sub-populations (i.e., gbMsM, underhoused persons, heterosexual persons, sex workers). Data from this section is also used to fulfill reporting requirements (i.e., sex and gender are both reported to federal partners).

Sexual contact information (pages 3-4 of [STI case investigation and contact tracing form](#)):

- Identified sexual partners, newborns or children of the case and their best available contact information.
- The *Notes* section should contain advice given with respect to specific control measures, specimens collected, date of collection, results and treatments administered. If the contact is identified as a case, specify this and initiate an STI case investigation form.
- Who to identify as a contact and the advice given will vary for the reportable disease as outlined in the respective [Communicable Disease Manual](#) chapter.

Where to find Reporting Forms

Case Investigation Form

Report STI cases and contacts to OCPHO utilizing the [STI case investigation form](#). Cases and/or contacts who cannot be followed-up after a [specific number of attempts](#) are to be reported to OCPHO using the [Syphilis & HIV Lost-to-Follow-up Report Form](#).

Contact Tracing Form

If additional space to document contact tracing information is required, the [STI Contact Tracing Form](#) can be used.

Important!

A [Case investigation and contact tracing form](#), even if not fully complete, must still be reported (submitted) to the OCPHO within the timeframes identified above. It is expected that HCPs submit an *updated* investigation and contact tracing form as new information is received. For example, it is unlikely that partner notification will be fully complete within the initial reporting timeframe but will be completed as per the Routine

Follow-up of STI Cases and Contacts chart and these updates need to be reported (submitted)

(<https://www.hss.gov.nt.ca/professionals/sites/professionals/files/resources/routine-follow-up-sti-cases-contacts.pdf>).

Completed report forms (initial forms and updates) should be sent to OCPHO by

Medical Confidential Fax: 867-873-0442

¹ See [Syphilis chapter of NWT Communicable Disease Control Manual](#)



SEXUALLY TRANSMITTED INFECTIONS CASE INVESTIGATION FORM

SECTION 1 – PATIENT INFORMATION

Infection(s) to report:	Chlamydia Trachomatis Syphilis	Gonorrhea	HIV ² (complete sections 1, 2, 6, and contact tracing form; contact OCPHO for additional reporting requirements)
--------------------------------	-----------------------------------	-----------	---

Affix Label	Last Name:	Penicillin allergy? Yes No Unknown Not asked
	First Name:	What is client's gender? Man Woman Other Not asked
	HCN:	Are the client's sexual partner(s) (select all that apply): Men Women Gender diverse Prefer not to answer Not asked
	Birthdate (dd/mmm/yyyy):	Is the client HIV positive? Yes No (<i>Negative test within the past year</i>) Unknown Prefer not to answer Not asked
	Sex at Birth: Male Female Intersex	
	Current Address:	Is the client currently pregnant? Yes; how many weeks: _____ No Unknown ³ Prefer not to answer Not asked

SECTION 2 – REASON FOR VISIT/TEST

Contact of a case Prenatal visit/screen Suspected gonococcal ⁴ or chlamydial ⁵ treatment failure	Test of cure Re-exposure or re-infection Self-reported sexual assault Symptomatic	Unprotected sex Routine screening Public health outreach Other: _____
--	--	--

SECTION 3 – HISTORY AND PHYSICAL EXAM – REQUIRED FOR SYPHILIS

A) HISTORY

Prior syphilis? Yes (date diagnosed): _____	No	Unknown	Prefer not to answer	Not asked
Rash Yes (mmm/yyyy): _____	No	Unknown	Prefer not to answer	Not asked
Chancre Yes (mmm/yyyy): _____	No	Unknown	Prefer not to answer	Not asked

B. PHYSICAL EXAM FINDINGS Patient declined physical exam

Oral (chancre, lesions): _____	Yes	No	Not examined
Skin (rash- general, palms, soles): _____	Yes	No	Not examined
Genital (chancre, lesions – please swab lesions): _____	Yes	No	Not examined
Perianal (chancre, lesions – please swab lesions): _____	Yes	No	Not examined
Lymph nodes (cervical, inguinal, generalized): _____	Yes	No	Not examined
Neurological (headache, vertigo, ataxia, ocular, personality changes): _____	Yes	No	Not examined

C. STAGING

Stage of syphilis: _____ Test pending Uncertain (consult ID specialist to complete staging)

SECTION 4 – TESTING INFORMATION

A) SPECIMEN COLLECTION – CHLAMYDIA & GONORRHEA

Date of collection (dd/mmm/yyyy): _____	Type of specimen: NAAT Culture (<i>charcoal swab</i>)
Collection site(s): Urethra Pharynx Urine Rectum Endocervix Eye Other (<i>specify</i>): _____	

² Additional information required. Physician or nurse practitioner to contact OCPHO Communicable Disease Unit for supplemental report form.

³ Do a pregnancy test if syphilis is suspected.

⁴ See sections 5-6 of [Canadian Guidelines on Sexually Transmitted Infections](#)

⁵ True antimicrobial resistance to *C. trachomatis* is rare. Consult OCPHO or an infectious disease (ID) specialist if repeat serologic response to treatment is inadequate.



B. SYPHILIS TESTING			
Syphilis serology taken: Yes No Patient refused	Swab of chancre/lesions? Yes Not offered Patient refused		
Requisition given No ability to test, reason: _____	N/A (no chancre or lesions present)		
POCT done? Yes No	Date of collection (dd/mmm/yyyy): _____		
Pregnancy test? N/A (male patient) No Yes, result: _____	Date (dd/mmm/yyyy): _____		

SECTION 5 - TREATMENT

A) CHLAMYDIA AND GONORRHEA	
Treatment Date (dd/mmm/yyyy): _____	Was patient empirically treated⁶? Yes No
Azithromycin 1gm PO	Ceftriaxone 250mg IM
Other (describe medication, dose, route, and frequency): _____	
Refused treatment	Treatment deferred; reason: _____

B) SYPHILIS *Consultation with a communicable disease specialist or colleague experienced in this area is recommended to assist with staging, treatment, and follow-up of infectious syphilis.*

Preferred treatment for all non-pregnant adults⁷ staged as primary, secondary or early latent syphilis (<1 year duration) is **Penicillin G benzathine (Bicillin L-A™) 2.4 million units deep IM injection as a single dose (DRUG NAME ALERT)**
 Alternate treatment if penicillin allergy (drug/dose/frequency): _____

Tx Date 1 (dd/mmm/yyyy): _____	Lot Number: _____	Was patient empirically⁶ treated? Yes No Treatment refused Treatment deferred; reason: _____
⁸Tx Date 2 (dd/mmm/yyyy): _____	Lot Number: _____	
⁸Tx Date 3 (dd/mmm/yyyy): _____	Lot Number: _____	

Serologic follow-up after treatment is required for all stages of syphilis⁷. Primary, secondary and early latent syphilis are to be monitored at 1, 3, 6 and 12 months after treatment.

SECTION 6 – RISK FACTORS

How many sex partners has the client had in the last 12 months? _____

Type(s) of sex: Oral sex Anal sex Vaginal sex Prefer not to answer Not asked

Does client use barriers (i.e. condom, dental dam) when you have sex? No Yes: How often? Always Sometimes Rarely

Where/how does the client report meeting sex partners?

Bars Online or dating apps Other, specify: _____
 At house parties or through friends On the street Prefer not to answer Not asked

Risk Factor	Yes	No	Prefer not to answer	Not asked
Has the client had any sex partners outside the NWT in the last 12 months?				
Has the client had any sex partners outside of Canada in the last 12 months?				
Has the client had a fixed (stable, permanent) address for the past 12 months?				
Has the client been incarcerated in the last 12 months?				
In the last year, has client <u>received</u> goods (i.e. drugs, shelter, money) in exchange for sex?				
In the last year, has client <u>given</u> goods (i.e. drugs, shelter, money) in exchange for sex?				

How often was the client's sexual decision-making compromised by alcohol or drugs in the last 12 months?

None of the time Some of the time Most of the time All of the time Prefer not to answer Unknown Not asked

What substance(s) have affected the client's sexual decision-making in the last 12 months (check all that apply):

Alcohol Injection drugs Crack cocaine Other, specify: _____ Prefer not to answer Not asked

⁶ Empiric treatment refers to antibiotic therapy that is administered prior to receipt of lab results that confirm disease diagnosis.

⁷ See [Canadian Guidelines on Sexually Transmitted Infections – Syphilis guide: Treatment and follow-up](#)

⁸ More than one dose may be required to sufficiently treat pregnant persons and those diagnosed with late latent syphilis. See [CDN Guidelines on STIs](#).



Are the client's partner(s) in the last 12 months (select all that apply):

Known

How many partners is the client not willing to provide contact information for? _____

Record contact information for all other known partners in contact tracing section.

Unknown/Anonymous

If contact is anonymous, record any descriptive information (i.e., gender, age, location, distinct physical characteristics, online/social media username or handle) in contact tracing section.

SECTION 7 – REPORTING INFORMATION

Office of the Chief Public Health Officer | Department of Health and Social Services Box 1320, Yellowknife NT X1A 2L9 | Phone: (867) 920-8646 | **Medical Confidential Fax: (867) 873-0442**

Completed by:

(Sign):

Phone:

Date (dd/mmm/yyyy):

Comments:

Report is: Initial Update **If update, new information provided on:** Syphilis staging Treatment Other



Report is: Initial Update

Index Case HCN:

MUST PRINT: SEXUALLY TRANSMITTED INFECTIONS (STI) CONTACT TRACING REPORT FORM⁹

Instructions: As per the [Reportable Disease Control Regulations](#) (R-128-2009), HCPs shall make reasonable attempts to initiate contact tracing within 24 hours of reportable disease diagnosis. Please submit initial available contact information with the case investigation form. HCPs are also to provide the OCPHO with information respecting the contact tracing and specific control measures that have been initiated or carried out. Please submit an updated contact tracing form whenever:

1. A new contact has been identified
2. A contact has been treated

SEXUAL CONTACT INFORMATION	
Contact Tracing	
Contact of a case of (check all that apply):	
Chlamydia	Gonorrhea
Syphilis	HIV
Affix Label	Last Name: _____
	First Name: _____
	HCN: _____
	Age: _____ Birthdate (dd/mmm/yyyy): _____
	Sex: Male Female Unknown Prefer not to answer
	Current Address: _____
	Confirmed Phone Number(s): _____
LAST EXPOSURE TO CASE Date (dd/mmm/yyyy): _____ Location (NWT community or out of territory): _____	
Relationship to case (check all that apply): Steady partner Ex-partner Casual unknown Casual known Sex worker	
Health care professional will follow-up with contact OR Contact information forwarded to _____ for follow up	
Follow up information	
Date contact notified (dd/mmm/yyyy): _____	
Date contact treated (dd/mmm/yyyy): _____ Specify drug, dosage, and route: _____	
If contact is lost-to-follow-up, report using Syphilis & HIV Lost-to-Follow-up Report Form	
Notes: 	
Report date (dd/mmm/yyyy): _____	Clinic name: _____
Report completed by (print): _____	Community: _____
Reported completed by (signature): _____	



Report is: Initial Update

Index Case HCN:

MUST PRINT: SEXUALLY TRANSMITTED INFECTIONS (STI) CONTACT TRACING REPORT FORM⁹

Instructions: As per the [Reportable Disease Control Regulations](#) (R-128-2009), HCPs shall make reasonable attempts to initiate contact tracing within 24 hours of reportable disease diagnosis. Please submit initial available contact information with the case investigation form. HCPs are also to provide the OCPHO with information respecting the contact tracing and specific control measures that have been initiated or carried out. Please submit an updated contact tracing form whenever:

1. A new contact has been identified
2. A contact has been treated

SEXUAL CONTACT INFORMATION				
Contact Tracing				
Contact of a case of (check all that apply):		Chlamydia	Gonorrhea Syphilis HIV	
Affix Label	Last Name:	LAST EXPOSURE TO CASE		
	First Name:	Date (dd/mmm/yyyy): _____ Location (community or out of territory): _____		
	HCN:	Relationship to case (check all that apply):		
	Age:	Birthdate (dd/mmm/yyyy):	Steady partner	Ex-partner Casual unknown
	Sex: Male Female Unknown Prefer not to answer		Casual known	Sex worker
	Current Address:			
Confirmed Phone Number(s):				
Health care professional will follow-up with contact OR Contact information forwarded to _____ for follow up				
Follow up information				
Date contact notified (dd/mmm/yyyy): _____				
Date contact treated (dd/mmm/yyyy): _____ Specify drug, dosage, and route: _____				
If contact is lost-to-follow-up, report using Syphilis & HIV Lost-to-Follow-up Report Form				
Notes:				
Report date (dd/mmm/yyyy):		Clinic name:	Community:	
Report completed by (print):		Reported completed by (signature):		

⁹ Contact Tracing Report Form is not to be scanned into Electronic Medical Record (EMR). Fax to the Office of the Chief Public Health Officer (OCPHO).



Report is: Initial Update

Index Case HCN:

MUST PRINT: SEXUALLY TRANSMITTED INFECTIONS (STI) CONTACT TRACING REPORT FORM⁹

Instructions: As per the [Reportable Disease Control Regulations](#) (R-128-2009), HCPs shall make reasonable attempts to initiate contact tracing within 24 hours of reportable disease diagnosis. Please submit initial available contact information with the case investigation form. HCPs are also to provide the OCPHO with information respecting the contact tracing and specific control measures that have been initiated or carried out. Please submit an updated contact tracing form whenever:

1. A new contact has been identified
2. A contact has been treated

SEXUAL CONTACT INFORMATION	
Contact Tracing	
Contact of a case of (check all that apply): Chlamydia Gonorrhea Syphilis HIV	
Affix Label	Last Name: LAST EXPOSURE TO CASE
	First Name: Date (dd/mmm/yyyy): _____ Location (NWT community or out of territory): _____
	HCN: Relationship to case (check all that apply):
	Age: Birthdate (dd/mmm/yyyy): Steady partner Ex-partner Casual unknown
	Sex: Male Female Unknown Prefer not to answer Casual known Sex worker
	Current Address:
Confirmed Phone Number(s):	
Health care professional will follow-up with contact OR Contact information forwarded to _____ for follow up	
Follow up information	
Date contact notified (dd/mmm/yyyy): _____	
Date contact treated (dd/mmm/yyyy): _____ Specify drug, dosage, and route: _____	
If contact is lost-to-follow-up, report using Syphilis & HIV Lost-to-Follow-up Report Form	
Notes:	
Report date (dd/mmm/yyyy):	Clinic name: Community:
Report completed by (print):	Reported completed by (signature):

⁹ Contact Tracing Report Form is not to be scanned into Electronic Medical Record (EMR). Fax to the Office of the Chief Public Health Officer (OCPHO).



Report is: Initial Update

Index Case HCN:

MUST PRINT: SEXUALLY TRANSMITTED INFECTIONS (STI) CONTACT TRACING REPORT FORM⁹

Instructions: As per the [Reportable Disease Control Regulations](#) (R-128-2009), HCPs shall make reasonable attempts to initiate contact tracing within 24 hours of reportable disease diagnosis. Please submit initial available contact information with the case investigation form. HCPs are also to provide the OCPHO with information respecting the contact tracing and specific control measures that have been initiated or carried out. Please submit an updated contact tracing form whenever:

1. A new contact has been identified
2. A contact has been treated

SEXUAL CONTACT INFORMATION	
Contact Tracing	
Contact of a case of (check all that apply): Chlamydia Gonorrhea Syphilis HIV	
Affix Label	Last Name:
	First Name:
	HCN:
	Age: Birthdate (dd/mmm/yyyy):
	Sex: Male Female Unknown Prefer not to answer
	Current Address:
Confirmed Phone Number(s):	
Health care professional will follow-up with contact OR Contact information forwarded to _____ for follow up	
Follow up information	
Date contact notified (dd/mmm/yyyy): _____	
Date contact treated (dd/mmm/yyyy): _____ Specify drug, dosage, and route: _____	
If contact is lost-to-follow-up, report using Syphilis & HIV Lost-to-Follow-up Report Form	
Notes:	
Report date (dd/mmm/yyyy):	Clinic name:
Report completed by (print):	Reported completed by (signature):

⁹ Contact Tracing Report Form is not to be scanned into Electronic Medical Record (EMR). Fax to the Office of the Chief Public Health Officer (OCPHO).